

Ravi Doctor DDS



Post Operative Instructions for Dental Implants

Here at Dr. Ravi Doctor's office, we care about our patients and we believe in making sure they have more than enough information to be able to take all the necessary precautions in order to make our procedures feel seamless. That is why we have come up with a list of post-operative instructions to help you take appropriate care of your new implants.

Please read below and pay special attention to our advice and suggestions:

Placement of dental implants usually does not create a great deal of pain. General post operative instructions apply, with the exception of rinsing.

Remember **DO NOT** rinse with the warm salt water. Dr. Doctor has given you a prescription mouth rinse containing chlorhexidine, an antibacterial medicine to assist in the healing and keep the site healthy.

Make sure that you get rid of any toothpaste residue prior to rinsing and then **DO NOT** eat or drink for 30 minutes after rinsing. This will maximize the potency of the medication.

Generally speaking following implant surgery the sutures need to be removed. If they become loose and annoying, call the office to have them removed at your earliest convenience.

1. Some degree of discomfort and pain arises as the numbness subsides. At the first sign of pain or discomfort, take the prescribed medications. As an alternative you may take 2-4 Advil. If you cannot take aspirin products, then take two Tylenol. If the pain you are having, is not controlled by the Advil or "Advil-like" prescription, you may take the prescribed severe pain medicine [typically Vicodin ES], but only after eating some food. Sometimes pain medications can cause nausea and vomiting. It is very important that you have some food in your stomach before you take them.

2. **DO NOT DISTURB THE AREA OF SURGERY.** Allowing the tissues to rest undisturbed assists the healing process. Avoid vigorous chewing, excessive spitting, or aggressive rinsing. If you routinely use a Water Pik, avoid doing so during the healing phase. Initial healing may be delayed, active bleeding restarted, or infection introduced.

3. Expect minor bleeding or oozing. If bleeding persists, continue biting on a fresh gauze with constant stable pressure for an additional 30 minutes to an hour. Biting on a moist teabag wrapped in gauze may help control persistent oozing from the surgical site. Tea contains tannic acid, which helps promote blood clotting.

Southlake Office: 261 East Southlake Blvd., #100 Southlake, TX 76092 817-328-2400

Arlington Office: 1810-A South Bowen Rd.

Arlington, TX 76013

817-274-8667

Ravi Doctor DDS



4. Firm pressure for 15-30 minutes usually controls the problem. If you check too frequently, (every few minutes), the blood clot will not form properly and bleeding will continue. Should active bleeding persist, please call the office or Dr. Doctor directly.

5. **LIMIT PHYSICAL ACTIVITY** during the first 24-48 hours after surgery. Over exertion may lead to postoperative bleeding and discomfort. When you lie down keep your head elevated on a pillow. You may wish to place a towel on your pillowcase to avoid staining from any blood-tainted saliva.

6. **PAIN FOLLOWING ORAL SURGERY** will be most severe within the first 6-8 hours after the operation. Please **DO NOT** drink alcoholic beverages while taking prescription medication. Do not wait for the pain to become unbearable before using some form of pain medication, as then it will be more difficult to control. Moderate to severe pain usually does not last longer than 48 hours. Mild discomfort usually diminishes after the third day. Please contact us if you experience prolonged severe pain so that we can help make you more comfortable.

7. **SWELLING RELATED TO THE SURGICAL PROCEDURE** may occur following certain types of procedures. An ice pack should be placed gently on the side of your face that the surgery has been performed on for 20 minutes, then taken off for 10 minutes and repeat steps for 1 hour. Anti-inflammatory medications, such as Advil, also help decrease swelling.

8. **FLUID INTAKE IS IMPORTANT.** Water, teas, broth, soups, or juices are all suitable. Avoid hot liquids until the numbness has worn off, and the bleeding has stopped.

9. **AVOID USING A STRAW FOR SEVERAL DAYS.** It may dislodge the blood clot and delay healing.

10. **FOOD SELECTION.** Soft, cool foods are most easily tolerated. A nutritious diet is most important to your comfort and temperament. Remember that eating can prevent nausea sometimes associated with certain medications. Supplements such as Ensure, Carnation Instant Breakfast and/or yogurt supply excellent added nutrition.

*** DO NOT EAT ANY POPCORN, POPPY SEEDS, SESAME SEEDS OR SEEDED FRUITS. THEY CAN GET STUCK AND CAUSE INFECTION!**

11. Take medications we have prescribed as directed. Yogurt with active cultures or acidophilus could be taken while on antibiotics to prevent diarrhea. It is important to take the antibiotics to completion. If you are given antibiotics and take birth control pills, you should be aware that the birth control may become ineffective, therefore take appropriate precautions.

Southlake Office: 261 East Southlake Blvd., #100 Southlake, TX 76092 817-328-2400

Arlington Office: 1810-A South Bowen Rd.

Arlington, TX 76013

817-274-8667

Ravi Doctor DDS



12. Take any regularly scheduled medication (for diabetes, high blood pressure, etc.) on your regular schedule unless advised to do otherwise.

13. TRY TO AVOID SMOKING COMPLETELY, as it tends to delay healing and interfere with regeneration.

The Day Following Surgery and Thereafter

- ANY SWELLING, SORENESS OR STIFFNESS IN THE JAW MUSCLES can be relieved by applying a warm moist towel to the affected side of the face.
- Sometimes a soft diet may be necessary for the first few days following surgery. Most patients are able to resume regular food intake within a short time.
- Bruising marks may appear on the skin of the face during the first few days after surgery. Should any undue reaction or complications arise, notify the office immediately.

If you need to contact us after office hours, please call the doctor's cell phone. If he does not answer immediately, remember to always leave a voice message with your name, number and where you can be reached.*

Dr. Doctor makes every attempt to answer his cell phone as promptly as possible. Occasionally, your messages do not reach him. If you are experiencing an emergency, go to the nearest hospital emergency room.

***If you have an emergency after hours you may call Dr. Doctor's cell: 817-233-6300**

Southlake Office: 261 East Southlake Blvd., #100 Southlake, TX 76092 817-328-2400

Arlington Office: 1810-A South Bowen Rd. Arlington, TX 76013 817-274-8667